

Lethal effects of pyrethrins on spruce budworm (*Choristoneura fumiferana*)

CAI Hong-jiao • YOU Min-sheng • FU Jian-wei • RYALL Krista • LI Shi-you

Received: 2009-09-13

Accepted: 2009-11-28

© Northeast Forestry University and Springer-Verlag Berlin Heidelberg 2010

Abstract: Spruce budworm (*Choristoneura fumiferana* (Clemens)) is one of the most serious forest insect pests in North America. Laboratory bioassays were performed to determine the lethal doses and lethal time of pyrethrins (a botanical insecticide) on 4th instar larvae of spruce budworm using larval dip assay. Results show that the LT_{50} values (time of 50% larval mortality) for spruce budworm at the pyrethrins concentrations of 12.5, 25, 50, 100, and 200 $\mu\text{g}\cdot\text{L}^{-1}$ were 94.78, 45.54, 20.36, 14.39 and 11.37 h, respectively. The percentage of cumulative mortality at the pyrethrins concentrations of 12.5, 25, 50, 100, 200 $\mu\text{g}\cdot\text{L}^{-1}$ was approximately 50%, 67%, 93%, 100% and 100% within 120 h, respectively. The LC_{50} value (concentration of 50% larval mortality) for the 4th instar larvae was 16.1 $\mu\text{g}\cdot\text{L}^{-1}$. Thus, larval mortality of spruce budworm increased in a concentration-dependent manner, and lethal time decreased with increasing pyrethrin concentrations. These findings suggest that pyrethrins have a potential in controlling spruce budworm populations.

Key words: Spruce budworm (*Choristoneura fumiferana*); botanical insecticide; pyrethrins

Foundation project: This research was supported by Natural Resources Canada, Canadian Forest Service. Project No. CFSCB2005-09.
The online version is available at [http:// www.springerlink.com](http://www.springerlink.com)

Cai Hong-Jiao
Fisheries College of Jimei University, Xiamen, Fujian 361021, P. R. China. Email: hongjiao@gmail.com

YOU Min-sheng (✉)
Institute of Applied Ecology, Fujian Agriculture and Forestry University, Fujian 350001, P. R. China. Email: msyou@fjau.edu.cn

FU Jian-Wei
Institute of Plant Protection, Fujian Academy of Agricultural Sciences, Fuzhou Fujian 350002, P. R. China

Ryall Krista
Canadian Forest Service, Natural Resources Canada, 1219 Queen Street East, Sault Ste. Marie, Ontario, P6A 2E5, Canada.

Li Shi-You
Canadian Forest Service, Natural Resources Canada, Building 57, 960 Carling Ave. Ottawa, Ontario, K1A 0C6, Canada.

Responsible editor: Zhu Hong

Introduction

The spruce budworm, *Choristoneura fumiferana* (Clemens), (Lepidoptera: Tortricidae), is one of the most destructive defoliators in the northeastern United States and eastern Canada (Cadogan et al. 2005). Its host plants include a variety of conifers (Talerico 1983), but spruce budworm prefers to feed on balsam fir (*Abies balsamea* [L.] Mill.), as well as spruce species (Cadogan et al. 2005). Chemical control was used to control spruce budworm outbreaks from 1950's to 1980's, such as synthetic chemicals of organophosphate, chlorinated hydrocarbons and carbamates (Nigam 1975; Shea and Nigam 1984).

The chemical pesticides can seriously threaten the endangered species, reduce biodiversity and induce the development of pesticide resistance and the resurgence of secondary pest populations. Thus, it is necessary to find the lower risk or environmentally-acceptable insecticides, based on entomopathogenic viruses, bacteria or fungi (EPA 2000). In particular, *Bacillus thuringiensis* subsp. *Kurstaki* Berliner (Btk) is one of the microbial insecticides that have been widely used successfully to control the spruce budworm in Canada since 1985 (Cadogan et al. 2004; Li et al. 2001). Although several types of viruses including nucleopolyhedrovirus (NPV), granulosis virus (GV), cytoplasmic polyhedrosis virus (CPV) and an entomopoxvirus (EPV) have been identified from spruce budworm populations (Cunningham 1985), field trials demonstrated that the efficacy of viruses was various (Cadogan et al. 2004). Hence, at present, the control of the spruce budworm over its range in North America is only dependent upon Bt-based insecticides. The repeated usage of Bt to control spruce budworm over a long time presents a great risk of pesticide resistance development on the spruce budworm. Thus, additional control options are practically needed.

The plant extracts such as rotenone from *Derris elliptica* (Ameen et al. 1983), azadirachtin from *Azadirachta indica* (Schmutterer 1981), capillin from *Artemisia nilagirica* (Banerji et al. 1990), arborine, a new bioactive compound related to quinazalone alkaloid, from *Glycosmis pentaphylla* (Muthukrish-

nan et al. 1999) and goniiothalamine from *Bryonopsis laciniosa* (Kabir et al. 2003) have all been demonstrated to have insecticidal properties for a variety of insect pest species. In this study, pyrethrin insecticides derived from pyrethrum from the Chrysanthemum flower (*Chrysanthemum cinerariifolium*) (Klaassen et al. 1996) were selected for testing against spruce budworm larvae. This pyrethrin insecticide not only provides a quick knockdown of insects and breaks down very quickly in the environment, but also has low risk of residual environmental contamination and very low toxicity to mammals (Hitmi et al. 2000). A pyrethroid is a synthetic chemical compound similar to the natural pyrethrins. Pyrethroid tends to be more effective than natural pyrethrins (Klaassen et al. 1996) and usually more stable in the environment than pyrethrins (Elliott 1989). Therefore, in the study, the botanical insecticide (natural pyrethrin) was tested to determine its lethal effect on 4th instar spruce budworm larvae.

Materials and methods

Insects

Newly-hatched 1st spruce budworm instar larvae were purchased from the Canadian Forest Service Insect Production Unit, Sault Ste. Marie, Ontario. Larvae were reared on artificial diet in 30-mL plastic cups at room temperature (approximately 24°C with 50%-60% R.H.). When larvae reached the 4th instar, they were selected as experimental insects.

Toxicity tests

Toxicity tests were conducted to determine the mortality of the budworm subjecting to pyrethrins 5.0 EC (Honghe senjun biology Co. LTD, P. R. China), containing 5% natural pyrethrum as the active ingredient. The pyrethrin 5.0 EC was diluted into six concentrations (200, 100, 50, 25, 12.5 and 6.25 $\mu\text{g}\cdot\text{L}^{-1}$) by using distilled water (pH = 6.8) based on preliminary mortality tests. The results of preliminary mortality tests showed that approximately 50% mortality resulted from 12.5 $\mu\text{g}\cdot\text{L}^{-1}$ pyrethrin treatment, whereas about 95% mortality resulted from 50 $\mu\text{g}\cdot\text{L}^{-1}$ pyrethrin treatment.

For each concentration, 30 randomly-selected larvae were immersed in the insecticide solution or distilled water (control) for three seconds. After the larvae were blotted dry on filter paper, they were reared in cups containing artificial diet and held at room temperature. Five larvae were reared in each cup. Each treatment was replicated three times (a total of 90 larvae/treatment). Larval mortality was recorded at 6, 12, 24, 48, 72, 96 and 120 h after treatment. Larvae were considered dead if they did not respond when lightly prodded with forceps.

Data analysis

The mean percentage of larval mortality was normalized by an arcsine-square-root and then subjected to one-way analysis of variance (ANOVA), followed by Duncan's test to compare dif-

ferences among the seven treatments at 0.05 significance level using DPS software (Tang and Feng 1997). However, only non-transformed means and variances are presented in Table 1. The LC₅₀ (concentration of 50% larval mortality) and LC₉₅ (concentration of 95% larval mortality) were estimated according to the probit analysis. The data on larval mortality over time were analyzed with a complementary log-log model (CLL) (Preisler and Robertson 1989; Robertson and Preisler 1992). The LT₅₀ (time of 50% larval mortality) and LT₉₅ (time of 95% larval mortality) were estimated according to the methods of Preisler and Robertson (1989). The lethal-dose ratio test was used to determine the differences between LT₅₀ and LT₉₅ at $\alpha = 0.05$ level (Robertson and Preisler 1992) using DPS software.

Results

Effects of different pyrethrin concentrations on mortality of spruce budworm larvae

Within the first 6 h after treatment, no significant difference ($p > 0.05$) of mortality of spruce budworm larvae was found at the pyrethrin concentrations of 50, 100, and 200 $\mu\text{g}\cdot\text{L}^{-1}$ (Table 1). No larvae died at the pyrethrin concentrations of 0.00 and 6.25 $\mu\text{g}\cdot\text{L}^{-1}$ within 12 h (Table 1). From 12 h to 72 h, a significant increase ($p \leq 0.05$) in mortality was found with increasing pyrethrin concentrations (Table 1). Mortality had a rapid increase within 24 h in all treatments. The cumulative percentage of larval mortality slowed down and reached a stable maximum value by day 4 (96 h). No mortality was found in the control group. The mortalities at the pyrethrin concentrations of 6.25, 12.5, 25, 50, 100, 200 $\mu\text{g}\cdot\text{L}^{-1}$ were about 5%, 50%, 67%, 93%, 100% and 100% at 120h after treatment respectively (Table 1). Overall, there was a significant increase in mortality with increasing pyrethrin concentrations (Fig. 1). When pyrethrins were at the concentrations of 100 and 200 $\mu\text{g}\cdot\text{L}^{-1}$, the 100% larval mortality occurred (Table 1). The rate of mortality at the pyrethrin concentration of 100 $\mu\text{g}\cdot\text{L}^{-1}$ was relatively slower than that in the 200- $\mu\text{g}\cdot\text{L}^{-1}$ pyrethrin concentration (Fig. 1).




Fig. 1 Cumulative mortality of spruce budworm larvae at 6, 12, 24, 48, 72, 96, and 120 h after treatments with different concentrations of pyrethrins 5.0 EC. The concentrations were 6.25, 12.5, 25, 50, 100, 200 $\mu\text{g}\cdot\text{L}^{-1}$. No larvae died in the control. The means of the three replicates \pm SE are shown.

Table 1. Percentage of cumulative mortality of 4th instar spruce budworm larvae treated with different concentrations of pyrethrins 5.0 EC in a larval dip assay

Concentration ($\mu\text{g}\cdot\text{L}^{-1}$)	Larval mortality \pm SE						
	6 h	12 h	24 h	48 h	72 h	96 h	120 h
0	0.00 \pm 0.00 c	0.00 \pm 0.00 f	0.00 \pm 0.00 g	0.00 \pm 0.00 g	0.00 \pm 0.00 g	0.00 \pm 0.00 f	0.00 \pm 0.00 f
6.25	0.00 \pm 0.00 c	0.00 \pm 0.00 f	2.22 \pm 1.92 f	2.22 \pm 1.93 f	2.22 \pm 1.93 f	3.33 \pm 3.33 e	4.44 \pm 1.93 e
12.5	0.00 \pm 0.00c	15.56 \pm 1.93 e	30.00 \pm 3.33 e	36.67 \pm 3.34 e	43.33 \pm 3.33 e	50.00 \pm 3.33 d	50.00 \pm 3.32 d
25	6.67 \pm 3.34 b	23.34 \pm 3.33 d	43.33 \pm 3.33 d	53.33 \pm 3.11 d	60.00 \pm 3.33 d	66.67 \pm 3.34 c	66.67 \pm 3.34 c
50	10.00 \pm 3.33 ab	30.00 \pm 3.33 c	65.56 \pm 5.09 c	76.67 \pm 3.34 c	86.67 \pm 3.34 c	93.33 \pm 3.34 b	93.33 \pm 3.34 b
100	13.33 \pm 3.34 a	43.33 \pm 3.34 b	78.86 \pm 5.10 b	90.00 \pm 3.21 b	92.22 \pm 1.93 b	97.78 \pm 1.91 a	100.00 \pm 0.00 a
200	13.34 \pm 3.34 a	56.67 \pm 3.22 a	88.89 \pm 1.93 a	97.78 \pm 1.93 a	100.00 \pm 0.00 a	100.00 \pm 0.00 a	100.00 \pm 0.00 a

Notes: 4th instar spruce budworm larvae (three replicates of 30 larvae per replicate) were tested using larval dip assay. The mortality was recorded on 6, 12, 24, 48, 72, 96, 120 h post-dip. The means of the three replicates \pm SE are shown. Means within columns followed by different letters were significantly different based on Duncan's multiple range test ($p \leq 0.05$).

Lethal dose and lethal time for spruce budworm larvae treated with pyrethrins

The LC_{50} was decreased with the time and was estimated as $23.91\mu\text{g}\cdot\text{L}^{-1}$ and $16.11\mu\text{g}\cdot\text{L}^{-1}$ in 48h and 120h, respectively (Table 2). The lethal time varied in a concentration – dependent manner. The LT_{50} values for spruce budworm at the pyrethrin concentration of 12.5, 25, 50, 100, and $200\mu\text{g}\cdot\text{L}^{-1}$ were estimated as 94.78, 45.54, 20.36, 14.39, and 11.37 h, respectively (Table 3). The LT_{95} values at the pyrethrin concentration of 50, 100 and $200\mu\text{g}\cdot\text{L}^{-1}$ were 119.71, 64.50 and 31.78 h, respectively (Table 3). Higher pyrethrin concentrations may cause larvae to die within shorter time periods (Table 3). The lethal-dose ratio test shows that there was a significant difference ($p \leq 0.05$) in LT_{50} between all pyrethrin concentrations, except for the concentration of 200 vs. $100\mu\text{g}\cdot\text{L}^{-1}$ (Table 4). LT_{95} ratio (95%CL) significantly increased with decreasing pyrethrin concentration except for the pyrethrin concentration of 12.50 vs. $25\mu\text{g}\cdot\text{L}^{-1}$ (Table 4). Thus, larval mortality of spruce budworm was increased in a concentration-dependent manner, and lethal time was decreased with increasing pyrethrin concentrations.

Table 2. Lethal concentrations for spruce budworm treated with different concentrations of pyrethrins 5.0 EC using larval-dip assay

Time (h)	Slope \pm SE	LC_{50} (95%CL) ($\mu\text{g}\cdot\text{L}^{-1}$)	χ^2	df
6	0.92 \pm 0.33	1978.84(341.61-535.48)	2.50	4
12	1.22 \pm 0.23	131.67(80.24-323.84)	2.98	4
24	1.79 \pm 0.24	33.62(25.29-44.33)	3.70	4
48	2.27 \pm 0.28	23.91(17.96-30.24)	3.96	4
72	2.55 \pm 0.33	19.82(14.48-25.05)	6.34	4
96	3.02 \pm 0.41	16.54(11.56-21.08)	5.34	4
120	3.23 \pm 0.44	16.11(11.17-20.51)	4.45	4

Notes: 4th instar spruce budworm larvae (three replicates of 30 larvae per replicate) were tested using larval dip assay. The mortality was recorded on 6, 12, 24, 48, 72, 96, 120h post-dip. The means of the three replicates \pm SE are shown. Each value of slope represented the regression slope of the relationship between larval mortality and lethal concentration.

Table 3. Lethal time for spruce budworm treated with different concentrations of pyrethrins 5.0 EC using larval dip assay

Pyrethrins ($\mu\text{g}\cdot\text{L}^{-1}$)	Slope \pm SE	LT_{50} (95%CL) (h)	LT_{95} (95%CL) (h)	χ^2	df
12.5	1.29 \pm 0.24	94.78 (63.70-191.70)	1770.89 (595.97-17793.5)	3.72	5
25	1.35 \pm 0.22	45.54 (33.38-65.89)	761.73 (343.83-3433.42)	2.01	5
50	2.14 \pm 0.26	20.36 (15.13-25.73)	119.71 (88.05-187.02)	1.67	5
100	2.52 \pm 0.30	14.39 (9.97-18.67)	64.50 (51.24-87.18)	2.70	5
200	3.69 \pm 0.54	11.37 (6.86-15.36)	31.78 (25.99-38.11)	0.68	5

Notes: 4th instar spruce budworm larvae (three replicates of 30 larvae per replicate) were tested using larval dip assay. The mortality was recorded at 6, 12, 24, 48, 72, 96, 120h post-dip. The means of the three replicates \pm SE are shown. Each value of slope represented the regression slope of the relationship between larval mortality and lethal time.

Discussion

The mortality of spruce budworm was higher after exposed to low concentrations of the pyrethroid and resmethrin (Morris 1975). However, the synthetic pyrethroid insecticides have disadvantages, including insecticide residues and potential human health risks. To mitigate these risks, pyrethroid insecticides were restricted from use in some areas. In 2006, EPA (Environmental protection agency in USA) announced the permethrin registration eligibility decision including application rates, droplet size and mandatory label statements.

In our preliminary study, higher mortality of spruce budworm was also found in all pyrethrin treatments in comparison with the control and the capsaicin-nicotine (botanical insecticide). Our results demonstrated that natural pyrethrins caused considerable mortality to larvae of the spruce budworm. Similar results were found in previous reports, demonstrating that pyrethrins have a considerable lethal effect on other insect species, such as leaf-

hopper (*Empoasca fabae*) (Maletta et al. 2006) and grape root borer (*Vitacea polistiformis*) (Weihman and Liburd 2006). In our study, the LC₅₀ of pyrethrins for spruce budworm was estimated

as 16.11 µg·L⁻¹. According to the study of Sheppard and Brad (2000), LC₅₀ of pyrethrin I and pyrethrin II for house flies (*Musca domestica*) was 0.20 µg·L⁻¹ and 0.49 µg·L⁻¹, respectively.

Table 4. Comparisons of lethal time for spruce budworm treated with different concentration of pyrethrins 5.0 EC using larval dip assay

Pyrethrins (µg·L ⁻¹)	LT ₅₀ ratio (95%CL)				
	12.5	25	50	100	200
12.5	1.00	2.08*(1.23-3.53)	4.66*(2.87-7.58)	6.59*(4.07-10.67)	8.34*(5.25-13.24)
25	0.50*(0.30-0.84)	1.00	2.24*(1.51-3.31)	3.17*(2.15-4.67)	4.01*(2.79-5.76)
50	0.21*(0.13-0.35)	0.43*(0.30-0.61)	1.00	1.42*(1.02-1.97)	1.79*(1.33-2.42)
100	0.15*(0.09-0.25)	0.30*(0.21-0.43)	0.71*(0.51-0.98)	1.00	1.27(0.94-1.70)
200	0.12*(0.08-0.19)	0.24*(0.17-0.33)	0.56*(0.41-0.75)	0.79(0.59-1.06)	1.00
Pyrethrins (µg·L ⁻¹)	LT ₉₅ ratio(95%CL)				
	12.5	25	50	100	200
12.5	1.00	2.33 (0.42-12.82)	14.85*(3.53-62.53)	56.06*(6.64-114.55)	56.06*(13.58-231.47)
25	0.31(0.06-1.48)	1.00	6.38*(2.17-18.76)	11.86*(4.11-34.20)	24.11*(8.43-68.97)
50	0.07*(0.02-0.28)	0.22*(0.09-0.51)	1.00	1.86*(1.10-3.14)	3.78*(2.27-6.28)
100	0.04*(0.01-0.15)	0.12*(0.05-0.27)	0.54*(0.32-0.91)	1.00	2.03*(1.27-3.25)
200	0.02*(0.01-0.07)	0.06*(0.03-0.13)	0.26*(0.16-0.44)	0.49*(0.31-0.79)	1.00

Notes: An asterisk (*) indicates a significant ($p \leq 0.05$) difference in LT₅₀ and LT₉₅ values between any concentrations of pyrethrins based on the 95% confidence limits (CL) of the ratio excluding 1 (Robertson and Preisler, 1992).

Benezet et al. (1988) found that the toxicity of the pyrethrins decreased with increasing temperature. As expected, there was a significant increase in mortality of spruce budworm with increasing concentrations from 12.5 to 200 µg·L⁻¹ from 24h to 96h. This result was in agreement with results of Barcic et al. (2006). Therefore, the spruce budworm larval mortality varied in a concentration-dependent manner. That is to say, the spruce budworm larval mortality is associated with time.

Previous studies have demonstrated that lethal time decreases with increasing concentrations of active ingredient (e.g. Li et al. 2003; Poland et al. 2006). Similarly, in our study, the lethal time decreased from approximately 120 h to 95 h with increasing of the concentration of pyrethrins. Appel et al. (2004) found that the LT₅₀ value of Drione (silica gel and synergized pyrethrins) decreased with increasing concentration and relative humidity in a study of the toxicity of Drione insecticidal against German cockroaches (*Blattella germanica*). Therefore, suitable humidity condition might accelerate the efficacy of pyrethrins at the same concentration, which needs further study.

This botanical insecticide is naturally derived from plants with low toxicity to mammals and fast biodegradation rate. Also, it has very low possible impact on development of insecticide resistance (Barcic et al. 2006). Based on our results, applications of low-dose (i.e., 50 µg·L⁻¹) pyrethrins could provide effective control for larval spruce budworm. Zapata et al. (2006) reported that the natural pyrethrins compounded piperonil butoxide (PBO) (LC₅₀ = 50.6 µg·L⁻¹ in 72 h) reduced the population of Mediterranean fruit fly (*Ceratitis capitata*). Igrc Barcic et al. (2006) reported that the application of pyrethrin with low dose (170 mL·ha⁻¹) significantly reduced leaf damage from the Colorado potato beetle (*Leptinotarsa decemlineata*). Therefore, pyrethrins may have potential to be used in controlling spruce budworm populations.

Acknowledgments

The authors greatly thank Prof. Huang Jin-Zhi for his help in revising the manuscript.

References

- Ameen M, Shahjahan RM, Khan HR, Chowdhury AKA. 1983. Toxicity of rotenone extracted from indigenous Derris roots on mosquito larvae. *J Bangladesh Acad Sci*, 7: 39-47.
- Appel AG, Gehret MJ, Tanley MJ. 2004. Effects of moisture on the toxicity of inorganic and organic insecticidal dust formulations to German cockroaches (Blattodea: Blattellidae). *J Econ Entomol*, 97: 1009–1016.
- Banerji A, Luthria DL, Kokate, SD. 1990. Toxicity of capillin, the insecticidal principle of *Artemisia nilagirica* Clarke. *Indian J Exp Bio*, 28: 588–589.
- Barcic JI, Bzaok R, Bezjak S, Culjak TG, Barcic J. 2006. Combinations of several insecticides used for integrated control of Colorado potato beetle (*Leptinotarsa decemlineata*, Say., Coleoptera: Chrysomelidae). *J Pest Sci*, 79: 223–232.
- Benezet HJ, Huffman BB, Helms CW. 1988. Comparative toxicity of selected insecticides to the cigarette beetle at different temperatures. *Tob Sci*, 32: 71–73.
- Cadogan BL, Scharbach RD, Brown KW, Ebling PM, Payne NJ, Krause RE. 2004. Experimental aerial application of a new isolate of the nucleopolyhedrovirus, CfmNPV against *Choristoneura fumiferana* (Lepidoptera: Tortricidae). *Crop prot*, 23: 1–9.
- Cadogan BL, Scharbach RD, Knowlesb KR, Krause RE. 2005. Efficacy evaluation of a reduced dosage of tebufenozide applied aerially to control spruce budworm (*Choristoneura fumiferana*). *Crop Pro*, 24: 567–553.
- Cunningham JC. 1985. Microbial Control of Spruce Budworms and Gypsy Moth. In: Grimble DG, Lewis FB. (ed.), *Proceedings Symposium*. Washington: Department of Agriculture, USA. pp. 61–67.
- Elliott M. 1989. The pyrethroids: early discovery, recent advances and the

- future. *Pestis Sci*, 27: 1337–351.
- EPA, 2000. *EPA strategic plan*. Washington DC, USA: United States Environment Protection Agency. pp. 1–77.
- Hitmi A, Coudret A, Barthomeuf C. 2000. The production of pyrethrins by plant cell and tissue cultures of *Chrysanthemum cinerariaefolium* and *Tagefes species*. *Critical Rev Plan Sci*, 19: 69–89.
- Igrc Baric J, Bazok R, Bezjak S, Gothlin Culjak T, Barcic J. 2006. Combinations of several insecticides used for integrated control of Colorado potato beetle (*Leptinotarsa decemlineata*, Say., Coleoptera: Chrysomelidae). *J Pest Sci*, 79: 223–232.
- Kabir KE, Khan AR, Mosaddik MA. 2003. Goniotalamin - a potent mosquito larvicide from *Bryonopsis laciniosa* L. *J Appl Entomol*, 127: 112–115.
- Klaassen CD, Amdur MO, Doull J. 1996. *Casarett & Doull's Toxicology - The Basic Science of Poison*. New York: McGraw-Hill, USA. pp. 35–68.
- Li Guiming, Zhang Xiangyue, Wang Luquan. 2001. The use of *Bacillus thuringiensis* on forest integrated pest management. *J Forestry Res*, 12(1): 51–54.
- Li SY, Skinner AC, Rideout T, Stone DM, Crummey H, Holloway G. 2003. Lethal and sublethal effects of a neem-based insecticide on balsam fir sawfly (Hymenoptera: Diprionidae). *J Econ Entomol*, 96: 35–42.
- Maletta M, Henninger M, Holmstrom K, Hort T. 2006. Potato leaf hopper control and plastic mulch culture in organic potato production. *HortTechnol*, 16: 199–204.
- Morris ON. 1975. Susceptibility of the spruce budworm, *Choristoneura fumiferana*, and the white-marked tussock moth, *Orgyia leucostigmata*, to *Bacillus thuringiensis*: chemical insecticide combinations. *J Inverteb Pathol*, 26: 193–198.
- Muthukrishnan J, Seifert K, Hoffmann KH, Lorenz MW. 1999. Inhibition of juvenile hormone biosynthesis in *Gryllus bimaculatus* by *Glycosmis pentaphylla* leaf compounds. *Phytochemistry*, 50: 249–54.
- Nigam PC. 1975. Chemical insecticides. In: Prebble RL (ed.), *Aerial Control of Forest Insects in Canada*. Ottawa: Department of the Environment, Canada. pp. 8–24.
- Poland TM, Haack RA, Petrice TR, Miller DL, Bauer LS. 2006. Laboratory Evaluation of the Toxicity of Systemic Insecticides for Control of *Anoplophora glabripennis* and *Plectrodera scalator* (Coleoptera: Cerambycidae). *J Econ Entomol*, 99(1): 85–93.
- Preisler HK, Robertson JL. 1989. Analysis of time dose-mortality data. *J Econ Entomol*, 82: 1534–1542.
- Robertson JL, Preisler HK. 1992. Experimental design of bioassays. In: Robertson JL, Russell RM, Preisler HK, Savin NE.(eds), *Pesticide bioassays with arthropods*. Boca Raton: CRC Press, FL. pp. 21–32.
- Schmutterer H. 1981. Some properties and components of the neem tree (*Azadirachta indica*) and their use in pest control in developing countries. *Meded Fac bouwwet-Rijksuniv Gent*, 46: 30–47.
- Shea PJ, Nigam PC. 1984. Chemical control. In: Schmitt DM, Grimble DG, Searcy JL (eds). *Spruce Budworm Handbook: Managing the Spruce Budworm in Eastern North America*. Washington: US Department of agriculture, USA. pp.115–132.
- Sheppard DC, Swedlund B. 2000. Toxicity of individual pyrethrin esters to house flies (Diptera: Muscidae). *J Entomol Sci*, 35(3): 279–282.
- Talerico RL. 1983. Summary of life history and hosts of the spruce budworms. In: Talerico RL, Montgomery M (eds). *Proceedings, Forest Defoliator—Host Interactions: A Comparison Between Gypsy Moth and Spruce Budworm*. Washington: US Department of Agriculture, Forest Service, USA. pp. 1–4.
- Tang Qingyi, Feng Mingguang. 1997. DPS data process system for practical statistics.. Beijing: China Science Press, pp. 300–450 (in Chinese).
- Weihman SW, Liburd OE. 2006. Mating disruption and attract-and-kill as reduced-risk strategies for control of grape root borer *Vitacea polistiformis*(Lepidoptera: sesiidae) in Florida vineyards. *Flo Entomol*, 89(2): 245–250.
- Zapata N, Budia F, Viñuela E, Medina P. 2006. Laboratory evaluation of natural pyrethrins, pymetrozine and triflumuron as alternatives to control *Ceratitis capitata adults*. *Phytoparasitica*, 34(4): 420–427.